

Prenatal DDT Exposure in Relation to Anthropometric and Pubertal Measures in Adolescent Males

**Beth C. Gladen, Mark A. Klebanoff, Mary L. Hediger,
Solomon H. Katz, Dana B. Barr, Mark D. Davis,
and Matthew P. Longnecker**
doi:10.1289/ehp.7287 (available at <http://dx.doi.org/>)
Online 7 September 2004



Prenatal DDT Exposure in Relation to Anthropometric and Pubertal Measures in
Adolescent Males

Beth C. Gladen (1), Mark A. Klebanoff (2), Mary L. Hediger (2), Solomon H. Katz (3),
Dana B. Barr (4), Mark D. Davis (4), Matthew P. Longnecker (1)

(1) National Institute of Environmental Health Sciences, National Institutes of Health,
Department of Health and Human Services

(2) National Institute of Child Health and Human Development, National Institutes of
Health, Department of Health and Human Services

(3) Krogman Center for Research in Child Growth and Development, University of
Pennsylvania

(4) National Center for Environmental Health, Centers for Disease Control and
Prevention, Department of Health and Human Services

correspondence to: Beth C. Gladen

mailing address: Biostatistics Branch, Mail Drop A3-03

National Institute of Environmental Health Sciences, P.O. Box 12233

Research Triangle Park, NC 27709, USA

package delivery: Room A352, David P. Rall Building, 111 T.W. Alexander Drive

phone: 919-541-3461

fax: 919-541-4311

e-mail: gladen@niehs.nih.gov

running title: Prenatal DDT and Anthropometry and Puberty in Boys

key words:

DDT

DDE

Growth

Puberty

Child development

Prenatal exposure delayed effects

Acknowledgements:

A conflict of interest was not reported.

abbreviations:

BMI: body mass index

CPP: Collaborative Perinatal Project

DHEAS: Dehydroepiandrosterone sulfate

ND: not detected

o,p'-DDT: 1,1,1-trichloro-2-(*o*-chlorophenyl)-2-(*p*-chlorophenyl)-ethane

p,p'-DDE: 1,1-dichloro-2,2-bis(*p*-chlorophenyl)-ethylene

p,p'-DDT: 1,1,1-trichloro-2,2-bis(*p*-chlorophenyl)-ethane

PBPP: Philadelphia Blood Pressure Project

SEI: socioeconomic index

Section Headers:

Abstract

Introduction

Materials and Methods

Results

Discussion

References

Table 1

Table 2

Table 3

Table 4

Table 5

Abstract

DDT, a pesticide once used widely in agriculture and now limited to public health use, remains a controversial chemical due to a combination of benefits and risks. DDT or its breakdown products are ubiquitous in the environment and in humans. Compounds in the DDT family have endocrine actions and have been associated with reproductive toxicity. A previous study reported associations between prenatal exposure to p,p'-DDE and increased height and weight in adolescent boys. We examined a group with higher exposures to see whether similar associations would occur. Our study group was 304 males born in Philadelphia in the early 1960s who had participated in a previous study. Anthropometric and pubertal measures from one to six visits during their adolescent years were available, as were stored maternal serum samples from pregnancy. We measured p,p'-DDE, p,p'-DDT and o,p'-DDT in the maternal serum. Outcomes examined in the boys were height, ratio of sitting height to height, body mass index, triceps skinfold thickness, ratio of subscapular to the sum of triceps and subscapular skinfold thicknesses, skeletal age, serum testosterone, and serum dehydroepiandrosterone sulfate. No associations between prenatal exposure to any of the DDT compounds and any outcome measure were seen.